



LIMESTONE
DENTISTRY

Office 812.288.9300
Fax 812.288.9602
Address 2948 E 10th Street Ste. A
Jeffersonville, Indiana 47130
Email om.limestonedentistry@gmail.com
Web limestonedentistry.com

Periodontal Scaling and Root Planing Post-Op Instructions

Scaling and root planing is a non-surgical treatment of periodontal disease. The purpose of the treatment is to remove bacterial plaque and tartar from around teeth and under the gum line, which is causing bone loss. The goal is to produce clean, healthy teeth and roots, which will promote healing of the inflammation and infection that causes gum disease.

You may take an over-the-counter pain reliever for any tenderness or discomfort. Take ibuprofen (Advil/Motrin) or acetaminophen (Tylenol) unless you have medical conditions or allergies.

After scaling and root planing, avoid eating anything on the area being treated for two hours or until the anesthetic has worn off completely. It is easy to bite or burn your cheek, tongue or lip while numb. Avoid any hard foods such as tortilla chips, potato chips, popcorn, or seeds for the next several days.

To help soothe the area, rinse your mouth 2-3 times a day with warm salt water rinses. If you are prescribed Peridex/Chlorhexidine, use as directed. Resume your home care regimen of brushing twice a day with a soft bristled toothbrush and daily flossing immediately, but be gentle with the area recently treated. Your gum health must be maintained with proper home care, as instructed, and regular dental visits. You may use a WaterPik if recommended.

It is not unusual for the teeth to be more sensitive to hot or cold temperatures, and/or sweets. This is normal. This occurs as the gum tissue heals and shrinks in size and should gradually resolve in a few weeks with proper home care. Consistently brushing two to three times daily with sensitivity toothpaste or using fluoride rinses may



LIMESTONE
DENTISTRY

Office 812.288.9300

Fax 812.288.9602

Address 2948 E 10th Street Ste. A

Jeffersonville, Indiana 47130

Email om.limestonedentistry@gmail.com

Web limestonedentistry.com

alleviate this over time. Avoid toothpastes with “whitening” or baking soda, as this will contribute to the problem.

If sensitivity continues or is severe, professional application of a desensitizing agent may be required.

Refrain from smoking for 24 to 48 hours after scaling and root planing as tobacco will delay healing of the tissues. Smoking cessation is highly recommended.

You can expect to notice less redness, less bleeding, and less swelling of your gum tissues. Your teeth may feel smoother and your mouth will taste and feel better.

We will look forward to seeing you for your regular periodontal maintenance every three to four months to keep up with what we just accomplished together.